


CBSE Test Paper 02

Chapter 08 Motion

1. Which of the following figures represent uniform motion of moving object correctly?

(1)



2. Which of the statement is true? (1)

Statement A: A curved line in a speed-time graph means non-uniform motion.


Statement B : A curved line in a velocity-time graph means uniform acceleration

- Statement A is false, B is true
- Both the statement A and B are true
- Neither statement A nor Statement B is true.
- Statement A is true, B is false

3. A body moving in a circle of radius r , covers $3/4^{\text{th}}$ of the circle. The ratio of the distance to displacement is **(1)**

- a. $3 : 2\sqrt{2}$
- b. $3\pi : 2\sqrt{2}$
- c. $3\sqrt{2} : 2\pi$
- d. $2\sqrt{2} : 3\pi$

4. The slope of the x-t graph is a measure of **(1)**



- a. velocity = 2ms^{-1}
- b. acceleration = 2ms^{-2}
- c. acceleration = $\frac{1}{2}\text{ms}^{-2}$
- d. velocity = $\frac{1}{2}\text{ms}^{-2}$

5. If a body starts from rest, what can be said about the acceleration of body? **(1)**

- a. Uniform accelerated
- b. Positively accelerated
- c. Negative accelerated
- d. Non-Uniform accelerated

6. The reference point from which the distance of a body is measured is called? **(1)**

7. How does the path of an object look graphically when it is in uniform motion? **(1)**

8. Which of the following is true for displacement? **(1)**

- a. It cannot be zero.
- b. Its magnitude is greater than the distance travelled by the object.

9. What is the quantity which is measured by the area occupied below the velocity-time

graph? **(1)**

10. Define one radian. **(1)**

11. The velocity of a body in motion is recorded every second as shown-

time (s)	0	1	2	3	4	5	6	7	8	9	10
velocity (m/s)	60	54	48	42	36	30	24	18	12	6	


Calculate the - **(3)**

- acceleration
- distance travelled and draw the graph.

12. An athlete completes one round of the circular track of diameter 200 m in 40 s. What will be the distance covered and the displacement at the end of 2 minutes 20 s? **(3)**

13. The velocity of a car is 18 ms^{-1} . Express this velocity in kmh^{-1} . **(3)**

14. The displacement - time graph for a body is given below. State whether the velocity and acceleration of the body in the region BC, CD, DE and EF are positive, negative or zero. **(5)**




15. Two stones are thrown vertically upwards simultaneously with their initial velocities u_1 and u_2 respectively. Prove that the heights reached by them would be in the ratio of $u_1^2 : u_2^2$ (Assume upward acceleration is $-g$ and downward acceleration to be $+g$). **(5)**


CBSE Test Paper 02
Chapter 08 Motion

Answer

1. a.



Explanation: Uniform motion of a moving object



2. d. Statement A is true, B is false

Explanation: When the speed of a body changes in an irregular manner, the speed-time graph of the body is a curved line. The distance traveled by the body is given by the area between the speed-time curve and the time axis. For uniform speed graph is straight line. For non-uniform speed graph is curved line. For Increasing speed graph Straight line sloping upwards. For decreasing speed graph Straight line sloping downwards.

When velocity – time graph is plotted for an object moving with uniform acceleration, the slope of the graph is a straight line.

So, statement A is true and B is false.

3. b. $3\pi : 2\sqrt{2}$

Explanation: $r' = r\sqrt{2}$ displacement

$$\text{distance} = \frac{3\pi r}{2}$$

$$\text{ratio} = \frac{3\pi r}{2}$$

$$\text{ratio} = 3\pi : 2\sqrt{2}$$

4. a. velocity = 2ms^{-1}

Explanation: The position-time graph is used in physics to describe the motion of an object over a period of time. Time, in seconds, is conventionally plotted on the x-axis and the position of the object, measured in meters, is plotted along the y-axis. The slope of the position-time graph reveals important information about the velocity of the object.

5. b. Positively accelerated

Explanation: If a body starts from rest, it starts moving. that means the change in velocity is positive. That means there is a POSITIVE acceleration.

6. The reference point from which the distance of a body is measured is called origin.

7. Ans. Graphically the path of an object will be linear i.e. look like a straight line when it is in uniform motion.

8. Both (a) and (b) are false with respect to concept of displacement.

9. The area occupied below the velocity-time graph measures the distance covered by any object.

10. It is the angle which is subtended at the centre by an arc having a length equal to the radius of the circle.

11. a. Acceleration = slope of the velocity time graph


$$a = \frac{V_2 - V_1}{t_2 - t_1}$$

$$a = \frac{54 - 24}{1 - 6} = \frac{30}{-5} = -6\text{m/s}^2$$


b. Distance = $S = ut + \frac{1}{2}at^2$

$$= 60 \times 10 + \frac{1}{2}(-6) \times (10)^2$$

$$= 600 - 300 = 300\text{m}$$



12. circumference of circular track = $2\pi r$
 $= 2 \times \frac{22}{7} \times \text{diameter} / 2$
 $= 2 \times \frac{22}{7} \times 200 / 2 = 4400/7 \text{ m}$
 rounds completed by athlete in 2min20sec = $s = 140/40 = 3.5$
 therefore, total distance covered = $4400 / 7 \times 3.5 = 2200 \text{ m}$



Since one complete round of circular track needs 40s so he will complete 3 rounds in 2mins and in next 20s he can complete half round therefore displacement = diameter = 200m.

13. Velocity = $\frac{18}{1000} \times \frac{1}{\frac{1}{3600}} = 64.8 \text{ kmh}^{-1}$

14. i. For AB, the curve is upward i.e. slope is increasing, therefore velocity is positive and remains same. So, $V = +ve$ but $a=0$
 ii. For BC, curve still has +ve slope so, $V = +ve$ but velocity is decreasing wrt time, therefore, $a = \text{negative}$
 iii. For CD, both velocity and acceleration are zero because there is no slope.
 iv. For DE, velocity v is increasing wrt time, so acceleration is +ve.
 v. For EF, velocity is +ve (positive slope of $x-t$ graph) but acceleration is zero because velocity remains same with time.

	AB	BC	CD	DE	EF
V	+ ve	+ ve	0	+ ve	+ ve
a	0	- ve	0	+ ve	0

15. By using velocity- distance equation, $v^2 - u^2 = 2aS$.

For upward motion, acceleration due to gravity, $a = -g \text{ m/s}^2$.

Let the height attained by object = 'h'

Therefore, We have, $v^2 = u^2 - 2gh$ or $h = \frac{u^2 - v^2}{2g}$ (1)

But at highest point $v = 0$

Therefore, equation (1) become; $h = \frac{u^2}{2g}$

Now, For first ball, Height attained, $h_1 = \frac{u_1^2}{2g}$ (2)

Similarly, For second ball, Height attained, $h_2 = \frac{u_2^2}{2g}$ (3)

Dividing equation, (2) by (3), we get.

$\frac{h_1}{h_2} = \frac{u_1^2/2g}{u_2^2/2g} = \frac{u_1^2}{u_2^2}$ or $h_1 : h_2 = u_1^2 : u_2^2$; Hence proved.

PE